Geometry Lomac 2015-2016		<b>Date</b> <u>1/6</u>	due <u>1/7</u>	Quadratics: Zeros	7.3L
Name LO:	I can identify the zeros of a qu	uadratic function.	Per	■ <b>P</b>	j


DO NOW On the back of this packet

## (1) Quadratics: Locating Zeros

The x-locations on any function where the output (the y-coordinate) is equal to zero are known, not surprisingly as the zeroes of the function. These are amazingly important in applied settings. When they are rational numbers then they can be found using a factoring technique. We'll develop the idea in the first exercise.

*Exercise* #1: Consider the quadratic function  $y = x^2 - 2x - 3$ . It's graph is shown below.

- (a) What are the zeroes of the function? Write their *x*-values and circle them on the graph.
- (b) Verify that the positive zero is correct by showing that y = 0.



(c) Factor the expression  $x^2-2x-3$ . How do these factors compare to the zeroes?

(d) Based on (c), determine where the zeroes of  $y = x^2 + 3x - 10$  are algebraically. Verify using a table. (THIS MEANS YOU MUST SHOW A SECTION OF THE TABLE THAT YOU USED.)

### (2) Quadratics: Zeros and the Zero Product Law

What is really going on here is perhaps the **second most important equation solving technique**, known as the **Zero Product Law**.

#### THE ZERO PRODUCT LAW

If two or more quantities have a product of **zero** then at least one of them must be equal to **zero**. In symbolic form:

If  $a \cdot b = 0$  then either a = 0 or b = 0 (or both are zero)

Exercise #2: Use the Zero Product Law to find all solutions to each of the following equations.

(a) 
$$(x+7)(x-2)=0$$

(b) 
$$(2x-1)(3x+4)=0$$

The **Zero Product Law** is remarkable because it allows us to solve equations with an  $x^2$  or higher level term in it, as long as the expression set equal to zero can be **factored**.

*Exercise* #3: Find the **roots** (solutions) to each of the following equations by using the **Zero Product Law**. Sometimes you will be instructed to **solve by factoring**.

(a) 
$$x^2 + 4x - 12 = 0$$


(b) 
$$2x^2 - 14x = 0$$

(c) 
$$x^2 - 25 = 0$$

(d) 
$$2x^2 + 5x - 12 = 0$$

# $\square$ (4) Completing the square

**Exercise** #4: Find the zeroes of the quadratic function  $y = 3x^2 - 6x - 24$  algebraically. Then verify your answer by using your calculator to sketch a graph of the parabola using the window indicated on the axes below. Clearly mark the zeroes on the graph.



## (4) Vertex form in applications

#### **APPLICATIONS**

5. A baking soda rocket is fired upwards with an initial speed of 80 feet per second. Its height, h, above the ground in feet can be modeled using the equation:

$$h(t) = -16t^2 + 80t$$
 where t is the time since launch in seconds

At what time, t > 0, does the rocket hit the ground? Find algebraically using factoring.

#### 

## REASONING

6. The two quadratic equations below have the same solutions. Can you determine why? Completely factor both to see what they have in common.

$$x^2 - 7x + 12 = 0$$

$$3x^2 - 21x + 36 = 0$$

(6) Exit Ticket

ON THE LAST PAGE

(7) Homework

**FLUENCY** 

- 1. The roots of  $x^2 6x 16 = 0$  can be found by factoring as
  - $(1) \{-16, 6\}$
- $(3) \{-2, 8\}$
- $(2) \{-8, 2\}$
- $(4) \{6, 16\}$
- 2. The equation (2x-3)(x+7) = 0 has a solution set of
  - (1)  $\left\{-7, 1\frac{1}{2}\right\}$
- $(3) \{-7, 3\}$
- (2)  $\{3, 7\}$
- $(4) \left\{ \frac{1}{2}, -3 \right\}$
- 3. Find the roots of each of the following equations by factoring:

(a) 
$$x^2 - 36 = 0$$

(b) 
$$x^2 + 12x + 27 = 0$$

(c) 
$$3x^2 + 5x - 2 = 0$$


(d) 
$$20x^2 - 10x = 0$$


(e) 
$$10x^2 + x - 21 = 0$$

(f) 
$$4x^2 - 16x - 84 = 0$$

The LO (Learning Outcomes) are written below your name on the front of this packet. Demonstrate your achievement of these outcomes by doing the following:

- 4. Consider the quadratic function  $y = x^2 4x 5$ .
  - (a) Using your calculator, graph the function on the grid provided.
  - (b) State the zeroes of the function by inspecting the graph. Circle their locations.





8			
DO NOW	Name	Date	Per

(1) Translation to algebra progress. Write one or more algebraic statement(s) to represent this situation. Be sure to write at least one "Let" statement to define any variables.

7.3L

Jay's father is twice as old as Jay. In 20 years Jay will be two-thirds as old as his father. How old is each now?